

DTIC

DTIC FILE COPY

2

SECURITY CLASSIFICATION OF THIS PAGE

SELECTED  
REPORT DOCUMENTATION PAGE

MAR 07 1990

Form Approved  
OMB No 0704-0188

1a REPORT SECURITY CLASSIFICATION (U)		1b RESTRICTIVE MARKINGS NA	
SECURITY CLASSIFICATION AUTHORITY NA		3 DISTRIBUTION/AVAILABILITY OF REPORT Distribution Unlimited	
DECLASSIFICATION/DOWNGRADING SCHEDULE NA			
PERFORMING ORGANIZATION REPORT NUMBER(S) Uniformed Services University of the Health Sciences		5 MONITORING ORGANIZATION REPORT NUMBER(S) NA	
NAME OF PERFORMING ORGANIZATION Uniformed Services Univ. of the Health Sciences		7a NAME OF MONITORING ORGANIZATION Office of Naval Research	
ADDRESS (City, State, and ZIP Code) Department of Surgery 4301 Jones Bridge Road Bethesda, Maryland 20814-4799		7b ADDRESS (City, State, and ZIP Code) 800 North Quincy Street Arlington, VA 22217-5000	
NAME OF FUNDING/SPONSORING ORGANIZATION Office of Naval Research		8b OFFICE SYMBOL (If applicable) ONR	
		9 PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER N0001488WM24020	
8c ADDRESS (City, State, and ZIP Code) 800 North Quincy Street Arlington, VA 22217-5000		10 SOURCE OF FUNDING NUMBERS	
		PROGRAM ELEMENT NO 61153N	PROJECT NO RR04108
		TASK NO 441f83301	WORK UNIT ACCESSION NO
11 TITLE (Include Security Classification) Cachectin/Tumor Necrosis Factor and the Pituitary-Adrenal Axis			
12 PERSONAL AUTHOR(S) Diana S. Malcolm, Ph.D.			
13a TYPE OF REPORT Final	13b TIME COVERED FROM 4/88 TO 12/89	14 DATE OF REPORT (Year, Month, Day) 90/2/28	5 PAGE COUNT 19
16 SUPPLEMENTARY NOTATION			
17 COSATI CODES		18 SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD 08	GROUP	Tumor Necrosis Factor, Cachectin, ACTH/Corticosterone thyroglobulin/macrophage/LPS/dexamethasone/cAMP	
SUB-GROUP			
19 ABSTRACT (Continue on reverse if necessary and identify by block number) Results from our studies investigating immune-neuroendocrine interactions have yielded interesting and novel findings. In unanesthetized rats, TNF (10ug/kg) was a potent stimulus for ACTH release without affecting hemodynamics. In vitro, TNF was without effect on the basal secretion of corticosterone (CS) but inhibited ACTH-stimulated CS release. Likewise, TNF inhibited TSH-stimulated thyroglobulin release from cultured human thyroid cells. In vitro, ACTH tended to inhibit LPS-induced TNF release from cultured macrophages, however, in the presence of serum-free media, ACTH appeared to potentiate TNF release. Also, the dose-response curves for CS and dexamethasone (inhibitors of TNF release) were shifted to the right. Together these findings suggest that factors present in sera and absent in serum-free media may interfere with LPS binding and/or uptake into macrophages. Results from these studies have provided additional knowledge and insight into the bidirectional communication between the neuroendocrine and immune systems.			
20 DISTRIBUTION/AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS		21 ABSTRACT SECURITY CLASSIFICATION (U)	
22a NAME OF RESPONSIBLE INDIVIDUAL Dr. Jeannine A. Najee		22b TELEPHONE (Include Area Code) (202) 696-4055	22c OFFICE SYMBOL ONR

DD Form 1473, JUN 86

Previous editions are obsolete

SECURITY CLASSIFICATION OF THIS PAGE

S/N 0102-LF-014-6603

90 03 05 064

AD-A218 837

## FINAL REPORT

### Cachectin/Tumor Necrosis Factor and the Pituitary-Adrenal Axis

ONR Contract No. N00014-88-WM-24020

#### Brief Statement of Objectives:

Several lines of evidence indicate that the immune system by way of macrophage hormones can influence neuroendocrine function and vice versa suggesting that these two systems are involved in a complete regulatory feedback loop. The purpose of our research studies was to determine whether the monokine TNF can alter pituitary ACTH and adrenal corticosterone release in vivo and in vitro and further to determine whether these neuroendocrine hormones can alter TNF release from macrophages in culture.

#### Specific Research Objectives:

1. Determine what effect Tumor Necrosis Factor (TNF) has on ACTH and corticosterone secretion in vivo in unanesthetized rats.
2. Determine what effect TNF has on the basal and stimulated release of ACTH and corticosterone from cultured pituitary and adrenal cells, respectively.
3. Determine whether ACTH and/or glucocorticoids (corticosterone, dexamethasone) can alter endotoxin-induced TNF release from cultured macrophages.

#### Research Results:

The first year of this project was devoted to the in vivo studies (Research Objective #1), as well as, a part of the in vitro studies involving cultured adrenal cells (Research Objective #2). Based on our observations of TNF's actions on adrenal cells, we also explored the effects of TNF on cultured human thyroid cells, specifically, the effects of TNF on TSH-induced thyroglobulin and cAMP production.

The second year of this project was largely devoted to the in vitro studies involving cultured macrophages (Research Objective #3). In addition to studying the effects of ACTH, corticosterone and dexamethasone on LPS-induced TNF release, we sought to determine whether these effects were altered when the macrophages were cultured with media (HL-1, Ventrex Corp) which was devoid of fetal calf serum.

We report here the results of our studies which are both interesting and new. In addition, results from these studies have provided possible insights into how the immune system by way of macrophage-derived peptides may regulate endocrine function, and conversely, how pituitary-adrenal hormones activated during infection or stress may act to regulate the release of TNF and thus modulate immune function.

1. Effects of recombinant human TNF-alpha on plasma levels of ACTH and corticosterone in the unanesthetized rat. (Figure 1 and 2)

TNF at doses of 0.01, 0.03 and 0.10 mg/kg was injected as a bolus into unanesthetized rats. Within 15 minutes of TNF injection, plasma ACTH levels were

maximal and not statistically different between the various doses of TNF administered. Thus, it appears that TNF induces a maximal ACTH response at 0.01 mg/kg which is not further elevated even with a ten-fold greater TNF dose (0.10 mg/kg). Although this dose of TNF did not produce significant changes in hemodynamics (data not shown) and was not associated with mortality, it appears to be a potent dose for neuroendocrine stimulation.

As expected, plasma corticosterone levels were elevated following TNF injection. This rise in plasma corticosterone is most likely ACTH-mediated, however, it is unknown from these in vivo studies alone whether TNF has direct effects on the adrenal gland to stimulate corticosterone release apart from or in addition to ACTH. Our in vitro experiments have attempted to answer some of these questions.

2. Effects of recombinant human TNF-alpha on basal and stimulated corticosterone release from cultured adrenal cells. (Table 1)

TNF alone at all doses tested (100, 300, 1000 ug/ml) had no effect on baseline corticosterone release from cultured adrenal cells. However, TNF clearly inhibited ACTH stimulated corticosterone secretion. This inhibition was reproducible and consistent and is seen at concentrations of TNF similar to that reported in patients with sepsis and with AIDS. There was no difference in cell number or viability following TNF application with or without ACTH present, indicating that the inhibitory effects of TNF are not due to cytotoxicity. This finding represents a significant and new interaction between the immune and endocrine systems.

3. Effects of recombinant human TNF-alpha on basal and stimulated thyroglobulin and cyclic AMP release from cultured thyroid cells (Tables 2 and 3).

Based on our observation of TNF's inhibitory actions on adrenal cells, we explored the effects of recombinant human TNF on human thyroid cells. TNF inhibited TSH stimulated thyroglobulin secretion from cultured thyroid cells in a dose-dependent manner. In all experiments, TSH exposure resulted in a brisk increase in cAMP production. However, even at the highest concentration, TNF had no effect on TSH stimulated cAMP production. This suggests that TNF's inhibition of TSH-stimulated thyroglobulin secretion is not mediated through cAMP.

4. Effects of ACTH on LPS-induced TNF release from cultured macrophages (Table 4 and Figure 3 & 4).

LPS (*E. coli* K235) at doses of 1, 10 and 100 ug x 10<sup>-4</sup> induced a biphasic dose-response effect on TNF release from macrophages cultured in the presence of Fetal Calf Serum (FCS) (Table 4). Maximal TNF release was elicited by the 10 ug x 10<sup>-4</sup> LPS dose, and this LPS dose was subsequently used to assess the effects of ACTH on stimulated TNF release from macrophages.

In the presence of FCS, ACTH appeared to inhibit LPS-stimulated TNF release (Table 4, Figure 3). However, judging from the 95% confidence interval associated with each mean value, there was no statistical difference between treatment groups. Since each value represents the mean of just 4-6 experiments, data from more experiments may show statistical significance between treatment groups.

Interestingly, macrophages cultured in HL-1 media (100x Concentrate; Ventrex Labs, Inc., Portland, ME) instead of FCS and challenged with LPS, exhibited a more linear dose-response in TNF release (Table 4). The biphasic response to LPS was not seen in macrophages cultured with HL-1.

In contrast to the observed effects of ACTH on macrophages cultured with FCS, in the presence of HL-1, ACTH appeared to potentiate LPS-stimulated TNF release (Figure 4). Because of the small sample size, statistical significance was not achieved, however, the trends in both treatment groups appear to be quite different suggesting that factors present in FCS may influence LPS-induced TNF release and its modulation by peptides.

5. Effects of Corticosterone on LPS-induced TNF release from cultured macrophages (Table 5)

As expected, corticosterone inhibited LPS-induced TNF release in a dose-related manner. Corticosterone's inhibitory effect appeared to be also present in the absence of FCS, however, owing to the small sample size this inhibitory effect was not statistically proven. The dose-response curve for corticosterone appeared to be shifted to the right in HL-1 cultured macrophages as compared to those cultured with 2% FCS.

6. Effects of Dexamethasone on LPS-induced TNF release from cultured macrophages (Table 6 and Figures 5 & 6)

Dexamethasone also inhibited LPS-stimulated TNF release from cultured macrophages, however, dexamethasone appeared to be a more potent TNF inhibitor than corticosterone when compared at the same molar concentrations (Tables 5 & 6). Dexamethasone's inhibitory effects were also present in the HL-1 cultured macrophages, however, the dose-response curve like that of corticosterone appeared to be shifted to the right in the presence of HL-1.

### Summary and Conclusions

Results from our studies investigating immune-neuroendocrine interactions have yielded several interesting and novel findings. In unanesthetized rats, TNF at a dose of 0.01 mg/kg was a potent stimulus for ACTH release. A dose of 0.1 mg/kg TNF did not further elevate this maximal ACTH response. Following TNF injection, plasma corticosterone was also elevated and most likely this release was mediated through ACTH since TNF was without effect on the release of corticosterone from cultured adrenal cells. At all doses tested, TNF was without significant effect on mean arterial pressure, heart rate or pulse pressure suggesting that TNF's effect on pituitary ACTH release was not secondary to a cardiovascular effect. Also our results indicate that TNF exerts a more potent effect on the pituitary-adrenal axis than on the cardiovascular system.

Interestingly, in vitro TNF inhibited ACTH-stimulated corticosterone release. It is unknown whether this inhibition also occurs in vivo, however, it is interesting to speculate that TNF may modulate the actions of ACTH in addition to influencing ACTH release. Glucocorticoids are potent inhibitors of TNF release both in vitro and in vivo and it is possible that TNF may influence corticosterone release (via ACTH) as part of a regulatory feedback loop.

TNF was also found to be a potent inhibitor of TSH-mediated thyroglobulin release from cultured human thyroid cells. The effects of TNF on the pituitary-thyroid axis are largely unknown. Clinically it has been observed that TNF is

elevated during medical conditions that are associated with the "sick euthyroid syndrome". One of the hallmarks of the "sick euthyroid syndrome" is the apparent suppression of thyroid hormone release in the face of adequate TSH stimulation. In other words, the thyroid gland's response to TSH is suppressed. It is interesting to speculate based on our observations that TNF may contribute to the thyroid gland suppression present in the "sick euthyroid syndrome" and warrants further study.

Data from our cultured macrophage studies suggest that ACTH may act to inhibit LPS-stimulated TNF release. If ACTH does influence macrophage secretion of TNF, this would imply that both the pituitary gland (ACTH) and adrenal gland (corticosterone) participate in the regulation of macrophage TNF release. This does not exclude the possibility that extra-pituitary sources of ACTH may likewise be involved. As expected, both corticosterone and dexamethasone potently inhibited TNF release in response to LPS.

Our studies using HL-1, a serum-free media, provided curious and new insights into the possible effects of serum factor(s) on LPS-induced TNF release. The LPS dose-response curve in macrophages cultured with serum-free media was linear and plateaued at the highest concentrations rather than being biphasic. The dose-response curves for both corticosterone and dexamethasone were shifted to the right compared to studies run using fetal calf serum in the tissue culture media. Most interesting was the observation that ACTH no longer appeared to inhibit but rather potentiated LPS-induced TNF release from macrophages cultured with serum-free media. These findings suggest that factor(s) present in sera and absent in HL-1 media may act to inhibit LPS-induced TNF release. It is unknown whether these factor(s) interfere with LPS binding and/or uptake into macrophages or whether they bind or restrict the activity of other modulators of TNF release (e.g. other peptides, hormones, calcium).

#### Publications

Poth, M., Tseng, Y.L., Wartofsky, L. Does Cachectin mediate altered thyroid function in systemic illness? A cell culture model. Presented at the 1989 Endocrine Society meeting.

Brennan, M.J., Betz, J.A., Poth, M. Tumor Necrosis Factor inhibits ACTH stimulated corticosterone secretion by rat adrenal cortical cells. Presented at the 1989 Endocrine Society meeting.

Malcolm, D.S., Poth, M. Tumor Necrosis Factor (TNF) and the pituitary-adrenal axis: In vivo and in vitro studies. Circulatory Shock (accepted).

Accession For	
NTIS - GRA&I	<input checked="" type="checkbox"/>
ERIC - TAB	<input type="checkbox"/>
Unpublished	<input type="checkbox"/>
Accession	
By	
Date	
Availability Codes	
Dist	Accession or Serial
A-1	

Figure 1. Effects of TNF on plasma levels of ACTH

## ACTH





Figure 2. Effects of TNF on plasma levels of corticosterone

# CORTICOSTERONE



**TABLE 1.** Effects of TNF on stimulated corticosterone release

Corticosterone secreted (ng/well)			
<u>ACTH</u>	<u>TNF (ug/ml)</u>	<u>Experiment I</u>	<u>Experiment II</u>
0	0	14 $\pm$ 2	6.4 $\pm$ 1.2
30	0	41.7 $\pm$ 7.8	16.9 $\pm$ 5.3
100	0	77.3 $\pm$ 6.2	63.0 $\pm$ 2.0
<hr/>			
30	100	2.6 $\pm$ .20	8.1 $\pm$ 5.3
30	300	.28 $\pm$ .03	
30	1000	.31 $\pm$ .02	3.6 $\pm$ .20
<hr/>			
100	100	1.21 $\pm$ .17	10.9 $\pm$ 1.7
100	300	3.98 $\pm$ 3.0	
100	1000	3.70 $\pm$ 2.15	3.5 $\pm$ 0
<hr/>			

Numbers are means of 3 wells

TABLE 2. Effects of TNF on basal and stimulated thyroglobulin release

Experiment I Thyroglobulin (ng/well) at 0-24 hours

	TNF (pg/ml)			
	0	100	300	1000
Without TSH	70 $\pm$ 6	79 $\pm$ 3	73 $\pm$ 1	65 $\pm$ 5
With TSH	340 $\pm$ 35	212 $\pm$ 25*	171 $\pm$ 17*	65 $\pm$ 21*

Experiment I Thyroglobulin (ng/well) at 24-48 hours

	TNF (pg/ml)			
	0	100	300	1000
Without TSH	114 $\pm$ 33	94 $\pm$ 10	79 $\pm$ 8	56 $\pm$ 12
With TSH	630 $\pm$ 126	251 $\pm$ 25*	160 $\pm$ 23*	63 $\pm$ 10*

Data is mean  $\pm$  S.D. using data from 3 wells

\*Different from control (TNF=0) with  $p < 0.01$

Experiment II Thyroglobulin (ng/well) at 24-48 hours

	TNF (pg/ml)			
	0	100	300	1000
Without TSH	369 $\pm$ 19	334 $\pm$ 5	258 $\pm$ 16*	223 $\pm$ 36*
With TSH	1025 $\pm$ 15	704 $\pm$ 44*	285 $\pm$ 51*	105 $\pm$ 11*

\*Data significantly different from control (TNF=0) according to ANOVA t-test ( $p < 0.05$ ).

**TABLE 3.** Effects of TNF on TSH-induced cAMP release

	<u>cAMP (picoles/well/2 hours)</u>			
	TNF (pg/ml)			
	0	100	300	1000
Without TSH	<0.5	<0.5	<0.5	<0.5
With TSH	5.6	5.0	4.7	5.1

---

Data are Mean of 3 wells.

Table 4 Effects of ACTH on LPS-induced TNF release in vitro

Treatment	TNF Bioactivity (U/ml)	
	Fetal Calf Serum*	HL-1 Media*
LPS (1 ug x 10 <sup>-4</sup> )	637 (265-1533)	160 (97-264)
LPS (10 ug x 10 <sup>-4</sup> )	11790 (6641-20931)	6741 (1947-23342)
LPS (100 ug x 10 <sup>-4</sup> )	6789 (4487-10270)	36098 (7421-175606)
LPS (500 ug x 10 <sup>-4</sup> )	--	81961 (34064-197205)
LPS (1000 ug x 10 <sup>-4</sup> )	--	96279 (46583-198988)
LPS (1 ug x 10 <sup>-4</sup> ) +		
--ACTH (10 <sup>-12</sup> M)	683 (183-2548)	--
--ACTH (10 <sup>-10</sup> M)	415 (175-982)	--
--ACTH (10 <sup>-8</sup> M)	537 (150-1923)	--
LPS (10 ug x 10 <sup>-4</sup> ) +		
--ACTH (10 <sup>-12</sup> M)	9082 (4836-17052)	6946 (1081-44623)
--ACTH (10 <sup>-10</sup> M)	3866 (925-16155)	10945 (3572-33557)
--ACTH (10 <sup>-8</sup> M)	11521 (6823-19458)	16059 (3874-66570)
LPS (100 ug x 10 <sup>-4</sup> ) +		
--ACTH (10 <sup>-12</sup> M)	--	59874 (8656-414157)
--ACTH (10 <sup>-10</sup> M)	--	129185 (47810-349060)
--ACTH (10 <sup>-8</sup> M)	--	144930 (66436-316159)
LPS (1000 ug x 10 <sup>-4</sup> ) +		
--ACTH (10 <sup>-12</sup> M)	--	42574 (2873-630961)
--ACTH (10 <sup>-10</sup> M)	--	>163898
--ACTH (10 <sup>-8</sup> M)	--	>163898

\* mean  $\pm$  95% confidence interval  
each value represents mean of 4-6 experiments

Figure 3. Effects of ACTH on LPS-induced TNF release from macrophages cultured in the presence of fetal calf serum. (Representative experiment)





Figure 4. Effects of ACTH on LPS-induced TNF release from macrophages cultured in the presence of HL-1 culture media. (Representative experiment)



**Table 5 Effects of Corticosterone (CS) on LPS-induced TNF release in vitro**

<u>Treatment</u>	<u>TNF Bioactivity (U/ml)</u>	
	<u>Fetal Calf Serum*</u>	<u>HL-1 Media*</u>
LPS (1 ug x 10 <sup>-4</sup> )	637 (265-1533)	160 (97-264)
LPS (10 ug x 10 <sup>-4</sup> )	11790 (6641-20931)	6741 (1947-23342)
LPS (100 ug x 10 <sup>-4</sup> )	6789 (4487-10270)	36098 (7421-175606)
LPS (500 ug x 10 <sup>-4</sup> )	--	81961 (34064-197205)
LPS (1000 ug x 10 <sup>-4</sup> )	--	96279 (46583-198988)
LPS (1 ug x 10 <sup>-4</sup> ) +		
--CS (10 <sup>-10</sup> M)	149 (99-225)	--
--CS (10 <sup>-8</sup> M)	117 (80-171)	--
--CS (10 <sup>-6</sup> M)	81 (79-82)	--
LPS (10 ug x 10 <sup>-4</sup> ) +		
--CS (10 <sup>-10</sup> M)	9009 (5409-10000)	14750 (1470-148004)
--CS (10 <sup>-8</sup> M)	2705 (1389-5265)	8417 (564-125492)
--CS (10 <sup>-6</sup> M)	356 (250-507)	637 (102-3987)
LPS (100 ug x 10 <sup>-4</sup> ) +		
--CS (10 <sup>-10</sup> M)	--	62944 (15741-251702)
--CS (10 <sup>-8</sup> M)	--	58395 (8242-413742)
--CS (10 <sup>-6</sup> M)	--	13440 (1498-120571)
LPS (1000 ug x 10 <sup>-4</sup> ) +		
--CS (10 <sup>-10</sup> M)	--	70404 (12991-381551)
--CS (10 <sup>-8</sup> M)	--	33962 (1458-790958)
--CS (10 <sup>-6</sup> M)	--	31225 (1133-860269)

\*mean  $\pm$  95% confidence interval

each value represents the mean of 4-6 experiments

**Table 6 Effects of Dexamethasone (DEX) on LPS-induced TNF release in vitro**

<u>Treatment</u>	<u>TNF Bioactivity (U/ml)</u>	
	<u>Fetal Calf Serum*</u>	<u>HL-1 Media*</u>
LPS (1 ug x 10 <sup>-4</sup> )	637 (26501533)	160 (97-264)
LPS (10 ug x 10 <sup>-4</sup> )	11790 (6641-20931)	6741 (1947-23342)
LPS (100 ug x 10 <sup>-4</sup> )	6789 (4487-10270)	36098 (7421-175606)
LPS (500 ug x 10 <sup>-4</sup> )	--	81961 (34064-197205)
LPS (1000 ug x 10 <sup>-4</sup> )	--	96279 (46583-198988)
LPS (1 ug x 10 <sup>-4</sup> ) +		
--DEX (10 <sup>-10</sup> M)	573 (143-2280)	--
--DEX (10 <sup>-8</sup> M)	98 (74-129)	--
--DEX (10 <sup>-6</sup> M)	<80	--
LPS (10 ug x 10 <sup>-4</sup> ) +		
--DEX (10 <sup>-10</sup> M)	9009 (5239-15490)	1194 (9-163570)
--DEX (10 <sup>-8</sup> M)	441 (169-1150)	226 (28-1806)
--DEX (10 <sup>-6</sup> M)	213 (116-389)	99 (65-153)
LPS (100 ug x 10 <sup>-4</sup> ) +		
--DEX (10 <sup>-10</sup> M)	--	7237 (904-57931)
--DEX (10 <sup>-8</sup> M)	--	1428 (217-9414)
--DEX (10 <sup>-6</sup> M)	--	371 (55-2500)

\*represents mean  $\pm$  95% confidence interval  
each value represents mean of 4-6 experiments

Figure 5. Effects of dexamethasone on LPS-induced TNF release from macrophages cultured in the presence of fetal calf serum (Representative experiment)





Figure 6. Effects of dexamethasone on LPS-induced TNF release from macrophages cultured with HL-1 media (serum free) (Representative Experiment)



DO NOT FOLD OR BEND THIS FORM. ABSTRACTS WILL BE PUBLISHED AS TYPED.

# The Endocrine Society

## 1989 ABSTRACT FORM

### IMPORTANT

Read all instructions before you type abstract. Also see sample abstracts and typing instructions on reverse side.

ALL ABSTRACTS MUST BE POSTMARKED

Outside USA — January 2, 1989

USA only — January 9, 1989

ONLY ACCEPTED ABSTRACTS  
WILL BE PUBLISHED.

### MAIL TO:

The Endocrine Society  
9650 Rockville Pike  
Bethesda, MD 20814  
Tel. (301) 571-1802

### PRESENTATION PREFERENCE

Do you strongly prefer a poster?

☐ Yes ☒ No

Preferences will not be binding on the  
Program Committee.

Indicate appropriate categories for your abstract  
(See Topic Category List):

1st Choice: #. 2. Title: Adrenal Cortex

2nd Choice: #. . . Title: . . . . .

Would you be willing to have your abstract taken  
out of its category and placed in a clinical session?

☒ Yes ☐ No

Do you think that your abstract is of interest to the  
lay public and, if asked, are you willing to write an  
abstract for the lay press?

☐ Yes ☒ No

1. The original typed copy of this abstract form (for  
reproduction by photo-offset in Program/Abstracts  
book) must be submitted together with 7 photocopies.

2. Your entire abstract, including a concise, descriptive  
title, author(s), sponsor if any, location where work was  
done, text and acknowledgments, must be within the  
rectangle outlined at the right. See reverse side for sample  
abstracts and full instructions.

3. Each Abstract Form submitted MUST BE SIGNED  
below by the member of THE ENDOCRINE SOCIETY.

Name and Telephone no. of presenting author

Michael J. Brennan

(202) 576-0055

TUMOR NECROSIS FACTOR INHIBITS ACTH STIMULATED CORTICOSTERONE SECRETION BY RAT ADRENAL CORTICAL CELLS. M.J. Brennan\*, J.A. Betz\*, and M. Poth. (SPON: D. Bunner) Walter Reed Army Medical Center, Washington, D.C. 20307 and The Uniformed Services University of the Health Sciences, Bethesda, M.D. 20814.

Tumor Necrosis factor (TNF) is a biologically active peptide secreted by macrophages and monocytes. TNF secretion is stimulated by endotoxin and TNF has been implicated in the pathogenesis of septic shock. To determine if TNF has specific actions on the adrenal gland, we studied the effects of ACTH and TNF on the *in vitro* secretion of corticosterone by rat adrenal cells. Adrenal glands from adult Sprague-Dawley rats were harvested, digested with collagenase, and cell suspensions were prepared. Cell cultures were incubated for 90 minutes in media with various concentrations of ACTH, TNF, or ACTH and TNF in combination. The cells were then centrifuged and the supernatants were assayed by RIA for corticosterone.

ACTH (pg/ml)	Corticosterone Secretion (pg/400,000 cells)		
	TNF (pg/ml)		
	0	100	1000
0	6.4 ± 1.2	5.8 ± 1.2	4.4 ± .20
30	16.9 ± 5.3	8.1 ± 5.3	3.6 ± .20
100	63.0 ± 2.0	10.9 ± 1.7	3.5 ± .00

Incubation with ACTH at concentrations of 10, 30 and 100 pg/ml produced a dose-response related stimulation of corticosterone secretion. Incubation with TNF alone at concentrations of 100, 300, and 1000 pg/ml had no effect on corticosterone secretion. However, when adrenal cells were incubated with ACTH and TNF in combination, corticosterone secretion was significantly inhibited. TNF at 100 pg/ml inhibited ACTH stimulated corticosterone secretion by 75-100% ( $p < .001$ ), while TNF at 1000 pg/ml produced 100% inhibition ( $p < .001$ ). Statistical significance was determined by multiple regression analysis. Conclusion: TNF inhibits ACTH stimulation of corticosterone secretion by rat adrenocortical cells. This is a new, potentially clinically important interaction between the immune and endocrine systems. Speculation: TNF may potentiate septic shock by inhibiting the body's ability to mount an appropriate glucocorticoid response to the stress of sepsis.

Member Signature Required Below as Sponsor of Abstract: NO MEMBER MAY SIGN MORE THAN ONE ABSTRACT.

The signing member certifies that any work with human or animal subjects related in this abstract complies with the guiding principles for experimental procedures as set forth in the Declaration of Helsinki, in the Statement of The Endocrine Society Concerning the Care and Use of Animals in Research, and in the NIH Guide for the Care and Use of Laboratory Animals, 1985. The signature also certifies that the scientific material in this abstract will not have been published or presented at any national meetings prior to The Endocrine Society Annual Meeting. Failure to adhere to this rule will result in deletion of the paper from the Program.

David L. Bunner  
Member signature

David L. Bunner  
Printed or typed name

(301) 663-7181  
Telephone number

# The Endocrine Society

## 1989 ABSTRACT FORM

**IMPORTANT**

Read all instructions before you type abstract. Also see sample abstracts and typing instructions on reverse side.

ALL ABSTRACTS MUST BE POSTMARKED

Outside USA — January 2, 1989

USA only — January 9, 1989

ONLY ACCEPTED ABSTRACTS  
WILL BE PUBLISHED.

**MAIL TO:**

The Endocrine Society  
9650 Rockville Pike  
Bethesda, MD 20814  
Tel. (301) 571-1802

**PRESENTATION PREFERENCE**

Do you strongly prefer a poster?

☐ Yes ☒ No

Preferences will not be binding on the  
Program Committee.

Indicate appropriate categories for your abstract  
(See Topic Category List):

1st Choice: #. 5. Title: Thyroid.....

2nd Choice: #. . . . . Title.....

Would you be willing to have your abstract taken  
out of its category and placed in a clinical session?

☒ Yes ☐ No

Do you think that your abstract is of interest to the  
lay public and, if asked, are you willing to write an  
abstract for the lay press?

☐ Yes ☒ No

1. The original typed copy of this abstract form (for  
reproduction by photo-offset in Program/Abstracts  
book) must be submitted together with 7 photocopies.

2. Your entire abstract, including a concise, descriptive  
title, author(s), sponsor if any, location where work was  
done, text and acknowledgments, must be within the  
rectangle outlined at the right. See reverse side for sample  
abstracts and full instructions.

3. Each Abstract Form submitted **MUST BE SIGNED**  
below by the member of THE ENDOCRINE SOCIETY.

Name and Telephone no. of presenting author

Dr. Merrily Poth

(202) 576-0055

DOES CACHECTIN MEDIATE ALTERED THYROID FUNCTION IN SYSTEMIC ILLNESS? A CELL CULTURE MODEL. M. Poth, Y.L. Tseng\*, and L. Wartofsky. Walter Reed Army Medical Center, Washington, DC 20307 and Uniformed Services University of the Health Sciences, Bethesda, MD 20814

Thyroidal economy in systemic non-thyroidal illness (SNTI) is marked by reductions in both central thyroid function and peripheral  $T_4$  to  $T_3$  conversion presumed to reflect a homeostatic mechanism to conserve energy. TSH levels tend to be normal in SNTI, and the mechanism underlying reduced thyroidal secretion is unknown. Recently, Ozawa (Endocrinol 123:1461, 1988) treated mice with tumor necrosis factor (TNF), as an animal model for SNTI, and reported diminished  $T_3$  and  $T_4$  responses to TSH administration. We have employed a primary thyroid cell culture system derived from surgical specimens to assess the effects of TNF on thyroidal responses to TSH. Cells at a density of 100,000/well were incubated with various concentrations (0-1000 pg/ml) of recombinant alpha-TNF (Genentech) and bTSH (1 mU/ml). Media were analyzed for cyclic AMP by RIA, and for thyroglobulin (Tg) by ELISA. TNF had no effect on either basal or TSH-stimulated cAMP generation.

Tg (ng/well) secreted into media by thyroid cells (100,000 cells/well) in the presence of TNF and bTSH (Mean  $\pm$  SEM)

TNF (pg/ml)	0-24 hr		24-48 hr	
	(-) TSH	(+) TSH	(-) TSH	(+) TSH
0	212 $\pm$ 7	365 $\pm$ 56	369 $\pm$ 19	1025 $\pm$ 15
100	186 $\pm$ 23	266 $\pm$ 57	334 $\pm$ 5	704 $\pm$ 44*
300	266 $\pm$ 73	144 $\pm$ 23*	258 $\pm$ 16*	285 $\pm$ 51*
1000	240 $\pm$ 41	89 $\pm$ 12*	223 $\pm$ 36*	105 $\pm$ 11*

\*Data significantly different from control (TNF=0) according to ANOVA t-test ( $p < 0.05$ ).

While TNF alone had no effect on Tg release at 24 hrs, TNF blunted TSH-stimulated Tg release by 27-76%. At 48 hrs, TNF blunted Tg release by 9-39% and TSH-stimulated Tg release by 31-90%. These results are consistent with the *in vivo* observations of Ozawa et al. and demonstrate a cytostatic effect of human thyrocytes by TNF in concentrations comparable to blood levels in man during SNTI. Thus, increases in circulating TNF in SNTI may be responsible for reduced thyroid function in these patients.

Member Signature Required Below as Sponsor of Abstract: NO MEMBER MAY SIGN MORE THAN ONE ABSTRACT.

The signing member certifies that any work with human or animal subjects related in this abstract complies with the guiding principles for experimental procedures as set forth in the Declaration of Helsinki, in the Statement of The Endocrine Society Concerning the Care and Use of Animals in Research, and in the NIH Guide for the Care and Use of Laboratory Animals, 1985. The signature also certifies that the scientific material in this abstract will not have been published or presented at any national meeting prior to The Endocrine Society Annual Meeting. Failure to adhere to this rule will result in deletion of the paper from the Program.

Dr. Merrily Poth

(202) 576-0055

Printed name

Telephone number

# THIRTEENTH ANNUAL CONFERENCE ON SHOCK

Tamarron

Durango, Colorado

June 8-11, 1990

H. Richard Adams, Scientific Program Chair

## ABSTRACT FORM

DEADLINE: Abstracts must be received by January 20, 1990

PLEASE READ CAREFULLY AND FOLLOW DETAILED INSTRUCTIONS PROVIDED ON REVERSE SIDE.

### NAME & MAILING ADDRESS OF FIRST AUTHOR

Dr. Diana S. Malcolm  
Department of Surgery  
Uniformed Services University  
of the Health Sciences  
4301 Jones Bridge Road  
Bethesda, Maryland 20814-4799

Telephone: (202) 295-3707

IMPORTANT: Please check if appropriate:

☐ Please send registration materials.

### PRESENTATION PREFERENCE (check one).

NOTE: See reverse side for details.

☒ Poster session ☐ Minisymposium ☐ Indifferent

### SELECT CATEGORY (See Topic Category List)

☒ TNF/Cytokines

☐ Hormones

Prepare abstract carefully on this form only. Return the original typed copy of this abstract plus 8 photocopies by January 26, 1990 to address on reverse side. Please enclose a non-refundable abstract handling fee of \$15.00 made payable to the Shock Society.

Abstracts will be reviewed by the Scientific Program Committee, H. Richard Adams, Chair.

TYPE ABSTRACT IN SPACE BELOW — DO NOT FOLD

**TUMOR NECROSIS FACTOR (TNF) AND THE PITUITARY-ADRENAL AXIS: IN VIVO AND IN VITRO STUDIES.** Diana Malcolm and Merrily Poth\*. Uniformed Services University of the Health Sciences, Bethesda, MD 20814.

Several lines of evidence indicate that the immune system by way of macrophage products can influence neuroendocrine function and vice versa suggesting that these two systems are involved in a complete regulatory feedback loop. The purpose of our studies was to determine what effect TNF has on ACTH and corticosterone (CS) release both in vitro and in vivo, and conversely, what effect ACTH and CS have on endotoxin-induced TNF release from cultured macrophages. Intravenous injections of low doses of TNF (0.01-0.10 mg/kg) in unanesthetized Sprague-Dawley rats (250-300 g) resulted in dose-related elevations in plasma ACTH and CS. In vitro, TNF (0, 0.1, 1.0 ng/ml) inhibited ACTH-induced CS release from cultured adrenal cells by 83% and 94%, respectively ( $p < 0.01$ ). TNF alone at these doses had no effect on the basal secretion of CS. Conversely, both ACTH ( $10^{-8}$  M- $10^{-12}$  M) and CS ( $10^{-6}$  M- $10^{-10}$  M) suppressed TNF release from endotoxin (LPS)-stimulated cultured macrophages in a dose-dependent manner. Furthermore, when macrophages were cultured in serum-free media (HL-1, Ventrex) instead of 2% fetal calf serum, the LPS dose-response curve was shifted to the right suggesting that factor(s) present in serum may facilitate LPS-induced TNF release. In summary, (1) plasma ACTH and CS levels are elevated following TNF injection, (2) TNF inhibits ACTH-stimulated CS release in vitro without affecting basal CS secretion and (3) both ACTH and CS inhibit TNF release from cultured macrophages. These findings represent new and significant interactions between TNF and the pituitary-adrenal axis and further support the existence of a regulatory feedback loop between the neuroendocrine and immune systems.

Paper to be judged for the Young Investigators Awards. My signature attests that this work will not be presented in any form at a national or international conference prior to the Shock conference.

NAME OF AUTHOR

Diana S. Malcolm, Ph.D.

*Diana S. Malcolm*

## DISTRIBUTION LIST

### Behavioral Immunology Program

Annual, Final and Technical Reports (1 copy each except as noted)

#### INVESTIGATORS

Dr. Itamar B. Abrass  
Department of Medicine  
University of Washington  
Harborview Medical Center  
Seattle, WA 98104

\* Dr. Christopher L. Coe  
Department of Psychology  
Harlow Primate Laboratory  
University of Wisconsin  
Madison, WI 53715

Dr. Prince K. Arora  
NIDDK, Bldg. 8, Room 111  
National Institutes of Health  
Bethesda, MD 20892

\* Dr. Sheldon Cohen  
Department of Psychology  
Carnegie-Mellon Univ.  
Pittsburgh, PA 15213

\* Dr. Andrew S. Baum  
Dept. of Medical Psychology  
Uniformed Services University  
of the Health Sciences  
4301 Jones Bridge Road  
Bethesda, MD 20814-4799

Dr. Walla L. Dempsey  
Dept. of Microbiology and  
Immunology  
Medical College of  
Pennsylvania  
3300 Henry Avenue  
Philadelphia, PA 19129

Dr. Charles A. Bowles  
Merrifield Research Lab, Inc.  
P.O. Box 2362  
Merrifield, VA 22116-2362

Dr. Robert L. Hunter  
Dept. of Pathology  
Emory Univ. School of  
Medicine  
WMB 760  
Atlanta, GA 30322

Dr. Karen Bulloch  
Dept. of Pediatrics M009D  
Univ. of California,  
San Diego, School of Medicine  
La Jolla, CA 92093.

Dr. Terry C. Johnson  
Division of Biology  
Ackert Hall  
Kansas State University  
Manhattan, KS 66506

Dr. Michael D. Cahalan  
Dept. of Physiology and  
Biophysics  
Univ. of California, Irvine  
Irvine, CA 92717

\* Dr. Jerome Kagan  
Dept. of Psychology  
Harvard University  
Cambridge, MA 02138

Dr. Donald A. Chambers  
801 S. Paulina - Room J30E  
Univ. of Illinois at Chicago  
P.O. Box 6998  
Chicago, IL 60680

\* Dr. Keith W. Kelley  
Lab. of Immunophysiology  
University of Illinois  
809 S. Wright Street  
Champaign, IL 61820-6219

Dr. James M. Krueger  
Dept. of Physiology  
University of Tennessee  
894 Union Street  
Memphis, TN 38163

- \* Dr. Sandra Levy  
WPIC  
Univ. of Pittsburgh  
School of Medicine  
3811 O'Hara Street  
Pittsburgh, PA 15213

Dr. Roger M. Loria  
Virginia Commonwealth Univ.  
Dept. of Microbiology &  
Immun. Box 678, MCV Station  
Richmond, VA 23298-0001

- \* Dr. Lester Luborsky  
Dept. of Psychiatry  
308 Piersol Bldg. G1  
Hospital of the University  
of Pennsylvania  
Philadelphia, PA 19104

- \* Dr. Steven F. Maier  
Dept. of Psychology  
University of Colorado  
Campus Box 345  
Boulder, CO 80309

Dr. Diana S. Malcolm  
Dept. of Surgery,  
Uniformed Services Univ.  
of the Health Sciences  
4301 Jones Bridge Road  
Bethesda, MD 20814-4799

Dr. Michael H. Melner  
Dept. of Reproductive Biology  
Oregon Regional Primate Center  
505 N.W. 185th Avenue  
Beaverton, OR 97006

Dr. Vera B. Morhenn  
Department of Dermatology  
Martinez VA  
Martinez, CA 94553

Dr. Jose R. Perez-Polo  
Dept. of Biochemistry  
Gail Borden Bldg., Room 436  
Univ. of Texas Medical Branch  
Galveston, TX 77550-2777

Dr. Merrily P.M. Poth  
Dept. of Pediatrics, A3027  
Uniformed Services Univ.  
of the Health Sciences  
4301 Jones Bridge Road  
Bethesda, MD 20814-4799

Dr. Eric M. Smith  
Dept. of Psychiatry  
Univ. of Texas Medical  
Branch  
Galveston, TX 77550

Dr. G. John Stanton  
Dept. of Microbiology  
Univ. of Texas Medical  
Branch  
Galveston, TX 77550

- \*\* Dr. Ross R. Vickers, Jr.  
Naval Health Research Center  
Building 346  
P.O. Box 85122  
San Diego, CA 92138

- \* Supported by ONR Code 1142BI
- \*\* Supported by NMRDC

Annual, Final and Technical Reports (one copy each except as noted)

ADMINISTRATORS

Dr. Jeannine A. Majde, Code 1141SB (2 copies)  
Program Manager, Systems Biology  
Office of Naval Research  
800 N. Quincy Street  
Arlington, VA 22217-5000

Program Manager  
Biotechnology Program  
Office of Naval Research  
Code 1213  
800 N. Quincy Street  
Arlington, VA 22217-5000

Administrator (2 copies) (Enclose DTIC Fm 50)  
Defense Technical Information Center  
Building 5, Cameron Station  
Alexandria, VA 22314

Program Manager  
Support Technology Directorate  
Office of Naval Technology  
Code 223  
800 N. Quincy Street  
Arlington, VA 22217-5000

Administrative Contracting Officer  
ONR Resident Representative  
(address varies - obtain from business office)

Mr. Michael McCracken  
218 3rd Ave NW  
Washington, DC 20006  
Commanding Officer  
Naval Medical Center  
Washington, DC 20372

DoD ACTIVITIES

Commander  
USAMRIID  
Fort Detrick  
Frederick, MD 21701

Commanding Officer  
Naval Medical Research & Development Command  
National Naval Medical Center  
Bethesda, MD 20814

Directorate of Life Sciences  
Air Force Ofc of Scient. Res.  
Bolling Air Force Base  
Washington, DC 20332

Director, Infectious Diseases Program Center  
Naval Medical Research Institute  
National Naval Medical Center  
Bethesda, MD 20814

Library  
Armed Forces Radiation  
Research Institute  
Bethesda, MD 20814-5145

Commander  
Chemical & Biological Sciences Division  
Army Research Office, P.O. Box 12211  
Research Triangle Park, NC 27709

Commander  
U.S. Army Research and Development Command  
Attn: SGRD-PLA  
Fort Detrick  
Frederick, MD 21701

Final and Technical Reports Only

Director, Naval Research Laboratory (6 copies)  
Attn: Technical Information Division, Code 2627  
Washington, DC 20375